板谷研究室 Itatani Group

研究テーマ Research Subjects

- 位相制御された高強度極短パルスレーザーの開発 Development of phase-stable intense ultrashort-pulse lasers
- 2 軟X線アト秒パルス発生と原子・分子・固体のアト秒分光 Generation of soft-X-ray attosecond pulse, attosecond spectroscopy of atoms, molecules, and solids
- 3 強レーザー場中での超高速現象の観測と制御 Measurement and control of ultrafast phenomena in strong optical fields
- 4 超高速軟 X 線分光法の開発 Development of ultrafast soft X-ray spectroscopy





専攻 Course 理学系物理学 Phys., Sci.

Professor ITATANI, Jiro




特任助教 水野 智也 特代 Project Research Associate Proj MIZUNO, Tomoya FUK



特任助教 深谷 亮 Project Research Associate FUKAYA, Ryo


高強度極短パルスレーザーの開発と、フェムト秒からアト秒領域の超高速現象に関する研究を行っている。光源開発に関しては、可視から中赤外領域での位相制御された高強度極短パルス光源の開発と、気体・固体・液体媒質での高次高調波発生を利用した多様な短波長パルス光源に関する研究を行っている。また、チタンサファイアレーザーを超えた次世代極短パルスレーザー光源を目指した光源技術の開発も進めている。光源利用に関しては、アト秒軟 X線パルスの超高速分光応用、原子・分子・固体中での高強度光電場で駆動された非線形光学現象に関する研究を主に行っている。位相制御された高強度極短パルス光源を基盤技術とした波長変換により、テラヘルツから軟 X 線までをカバーした超高速分光が実現可能であり、物質の非平衡状態における動的過程を様々な自由度を通して実時間観測し、さらには光で制御することを目指している。

We are working on the development of intense ultrashort pulse light sources and their applications in ultrafast spectroscopy on the femtosecond to attosecond time scale. In light source R&D, we focus on the generation of waveform-controlled intense optical pulses from the visible to the mid-infrared spectral range and the generation of shortwavelength ultrashort pulses using the physics of high-order harmonic generation in gases, solids and liquids. In addition, we are developing the building blocks of next-generation light sources to overcome the limitations of current Ti:sapphire laser-based technologies. Based on these novel light sources and techniques, we are developing attosecond soft X-ray spectroscopy and other ultrafast methods to probe fielddriven nonlinear processes in atoms, molecules, solids, and liquids. Our waveform-controlled intense light sources and related technologies will enable novel ultrafast spectroscopy covering an extremely broad spectral range from THz to soft x-rays. Our goal is to observe and control the ultrafast dynamics of non-equilibrium states of matter through multiple degrees of freedom.



アト秒軟 X線パルスを用いた N_2O 分子の過渡吸収分光と、内殻励起に関与するエネルギー準位、観測されたサブサイクルの変調を受けた過渡吸収スペクトル。

Schematic of transient absorption spectroscopy of N_2O molecule using attosecond soft x-ray pulses, energy levels involved in inner-shell excitation, and the observed transient absorption spectra. The observed ultrafast modulation is due to the tunnel ionization of the molecule in a core-hole state.



(左)強レーザー場中での光電子の再散乱によって得られるキャリア・エンベロープ位相依存性に依存した光電子スペクトル、(右)観測された光電子スペクトルから再構成された微分散乱断面積と理論との比較。

(Left) Carrier-envelope phase dependence of the photoelectron spectra observed by rescattering of laser-accelerated photoelectrons. (Right) Comparison of the differential scattering cross section reconstructed from the observed phase-dependent photoelectron spectra. The good agreement indicates that quantitative information can be obtained from the high-energy rescattering phenomena.

https://www.issp.u-tokyo.ac.jp/maincontents/organization/labs/itatani_group.html