

近藤研究室 Kondo Group

研究テーマ Research Subjects

- 1 極限レーザーを励起光源とする超高分解能角度分解光電子分光装置の開発
Development of a laser-excited ARPES system with ultra-high energy resolution
- 2 角度分解・スピン分解・時間分解光電子分光で見る超伝導やトポロジカル量子相
Superconductivity and topological quantum phase investigated by angle-, spin-, and time-resolved photoemission spectroscopy
- 3 放射光を利用した光電子分光で研究する強相関電子系物理
Strongly correlated physics studied by photoemission with synchrotron radiation




准教授 近藤 猛
Associate Professor KONDO, Takeshi

専攻 Course
理学系物理学
Phys., Sci.

固体中の電子が描くバンド構造は、あらゆる電子物性を理解する上での基礎を与える。角度分解光電子分光は、光で励起する光電子を角度及びエネルギーの関数としてイメージングすることでバンド構造を可視化する強力な実験手法である。この技術をベースとして、バンドが持つスピン構造を同定したり、パルス光で制御する非平衡ダイナミクスをフェムト秒スケールで観測することで、多彩な電子物性がバンド構造を通して見えてくる。当研究室では、このような卓越した光電子分光技術を駆使して、(高温)超伝導体、重い電子系や電子相関系物質、トポロジカル量子相、固体表面や薄膜で制御する量子井戸構造などの電気磁気物性を、直接観察で得られるバンド構造を舞台に研究する。さらには、極限レーザー光源及びそれを用いた高精度な光電子分光装置を開発し、フェルミ準位極近傍の微細な電子構造（エネルギーギャップや素励起との相互作用）を解明する。

The momentum-resolved band structure provides fundamental information to understand the electronic properties of materials. The angle-resolved photoemission spectroscopy (ARPES) is a powerful technique to visualize the band structure by mapping the intensities of photoelectrons as a function of angle and energy. With the spin-resolved technique, we can also identify the spin-polarized character of the band. In addition, the time-resolved ARPES realized with a pump-probe technique can track the reordering process of electron system from its nonequilibrium state. In our laboratory, we utilize these various ARPES techniques and study the following phenomena: nonconventional superconductors, heavy fermions, strongly correlated systems, topological quantum phases, and quantum well states. Furthermore, we develop a new ARPES machine capable of achieving both the lowest measurement temperature and the highest energy resolution in the world by innovating a ^3He cryostat and a laser source. The state-of-art equipment will enable us to identify even a subtle electronic feature close to the Fermi level, such as an energy gap and a mode-coupled dispersion, which is typically tied to exotic behaviors of conduction electrons.



(a) 銅酸化物高温超伝導体 $\text{Bi}_2\text{Sr}_2\text{CuO}_{6+d}$ の結晶構造。(b) 光電子アナライザー。(c) 角度分解光電子分光実験の模式図。(d) バンド分散のスナップショット。(e) バンド構造の全体図。(f) フェルミ面周りで描く超伝導と擬ギャップの競合関係。(g) 超伝導転移温度 (T_c) より高温 (黒線) と低温 (赤線) で測定したフェルミ面周りのスペクトル。(h) (g) で示すスペクトルの T_c 上下での差分。超伝導成分が赤で塗られており、(f) で模式的に示す赤の領域と対応する。

(a) Crystal structure of $\text{Bi}_2\text{Sr}_2\text{CuO}_{6+d}$ high- T_c superconductor. (b) ARPES analyzer. (c) Diagram of ARPES experiment. (d) Snapshot of dispersion image. (e) Whole band structure. (f) Competition between superconducting gap and pseudogap. (g) Spectra around Fermi surface below (red) and above (black) superconducting transition temperature ($T_c = 35\text{K}$). (h) Difference between the curves in (g). Coherent spectral weight is painted with a red color, which is corresponding to the red region represented in (f).



https://www.issp.u-tokyo.ac.jp/maincontents/organization/labs/kondo_group.html