

Spin and nuclear spin dependent transport in quantum dots

Keiji Ono¹ and Seigo Tarucha^{1,2}

¹*Dept. of Physics, University of Tokyo, 7-3-1 Hongo Bunkyo-ku, Tokyo 113-0033, Japan*

²*ERATO Mesoscopic Correlation Project, JST*

ono@phys.s.u-tokyo.ac.jp

1. Introduction

Zero-dimensional electronic system can be confined in semiconductor quantum dots [1]. The electronic state in the quantum dot, including the spin configuration, can be well controlled as a function of plunger gate voltage and magnetic field [2]. Recently, spin and nuclear spin degrees of freedoms in quantum dot are subject to intensive studies from the viewpoints of applications to future spintronics and quantum computations [3]. This paper presents our recent studies on the electron transport properties of quantum dots, concerning how the spins and nuclear spins in the quantum dots can influence the electron transport.

2. Device

We use two quantum dots connected in series between source/drain contacts, which are incorporated in a gated sub-micron pillar of triple barrier resonant tunneling structure: 8.0nm (6.0nm) thick $\text{Al}_{0.22}\text{Ga}_{0.78}\text{As}$ outer (center) barriers and 12nm thick two $\text{In}_{0.05}\text{Ga}_{0.95}\text{As}$ wells (Fig. 1) [4]. A DC magnetic field is applied parallel to the wells. An AC magnetic field of frequencies up to 15MHz, which is generated by a coil near the device, is applied perpendicular to the wells.

3. Spin Pauli blockade [5]

When the electrostatic potentials for two dots are relevantly adjusted, three charge states $(N_1, N_2) = (0,1)$, $(1,1)$ and $(0,2)$, where N_1 (N_2) is the number of electrons in dot 1 (dot 2), are degenerate and at the same time Coulomb blockade is lifted at zero source drain voltage V_S . The $(1,1)$ state contains two spin-states, a singlet and a triplet, with a slight energy difference by the amount of inter-dot exchange energy (estimated to be $\sim 100\mu\text{eV}$ or less). On the other hand the $(0,2)$ state only takes a spin singlet because the two electrons share the same orbital of the dot 2 due to Pauli exclusion principle. By applying negative V_S (Fig. 2(a)) current is carried by the cycle of three irreversible steps as $(0,1) \rightarrow (0,2)$ singlet $\rightarrow (1,1)$ singlet $\rightarrow (0,1) \dots$. For positive V_S (Fig. 2(c) and (f)), however, although there is a current carrying cycle of $(0,1) \rightarrow (1,1)$ singlet $\rightarrow (0,2)$ singlet $\rightarrow (0,1) \dots$, once the other transition $(0,1) \rightarrow (1,1)$ triplet takes place, further electron transfer is prohibited by Coulomb blockade and Pauli exclusion.

Figure 3 shows a current voltage characteristic at certain V_G where a large conductance peak first appeared at zero bias following the scheme of Fig. 2(b). For V_S larger than 1mV, the transport is in the nonlinear regime and the current is suppressed. This suppression is lifted for $V_S > 7\text{mV}$, where the Fermi energy of the right reservoir is lower

than $N_2=1$ state of the right dot.

4. Leakage current in spin blockade regime

In this spin blockade state a small leakage current of order $\sim 1\text{pA}$ has been observed. This means the $(1,1)$ triplet has a finite lifetime of order $e/I \sim 100\text{ns}$ (e , elementary charge). Possible causes for the leakage current are spin scattering that changes the state from $(1,1)$ triplet to $(1,1)$ singlet and co-tunneling processes that can directly scatter $(1,1)$ triplet to $(0,1)$ [5].

The leakage current in this spin blockade region is measured as a function of DC magnetic field. When the magnetic field is increased, the current increases rapidly at 0.5T and starts to oscillate as a function of time at about 0.7T. Both of the period and amplitude of the current oscillations increase with magnetic field, and disappear at about 0.9T. Figure 4(a) shows the data taken at 0.85T. A strong reduction in the oscillation period and amplitude of the current is observed when AC magnetic field whose frequency matches to ^{71}Ga nuclear spin resonance, is applied (Fig. 4(b)). This NMR-like behavior is only observed for the spin blockade region.

The origin of the current oscillations and the NMR response are not yet understood. Here we only outline our tentative explanation below. The spin scattering from $(1,1)$ triplet to $(1,1)$ singlet can be possible via the hyperfine flip-flop scattering with nuclei in quantum dots when the energy differences of the two spin states (electron spins and nuclear spins) are compensated [7]. This compensation will be achieved at a certain magnetic field where one of the Zeeman-split $(1,1)$ triplet states is degenerate with $(1,1)$ singlet. Then the hyperfine flip-flop scattering provides the nuclei with the same spin momentum, i.e., “down” nuclear spin can be flopped to “up” but not for vice versa. Because of the long relaxation time of nuclear spin at low temperature (up to few tens of min), the flopped nuclear spins can be accumulated during many cycles (period $\sim 100\text{ns}$) of the $(1,1)$ triplet occupation followed by scattering from $(1,1)$ triplet to $(1,1)$ singlet. This eventually leads to dynamical polarization of nuclei, which can acts back to the electronic state and hence the electron transport through the two-dot system via the Overhauser effect [8].

Acknowledgements

This work is supported from the Specially Promoted Research, Grant-in-Aid for Scientific Research, from the Ministry of Education, Science and Culture in Japan and from CREST-JST.

References

[1] L.P. Kouwenhoven et.al., in: *Electron transport in quantum*

dots, in *Mesoscopic Electron Transport*, edited by L.L. Sohn, G. Schon, L.P. Kouwenhoven (Kluwer, Series E345, 1997).

[2] S. Tarucha, et.al, Phys. Rev. Lett. **77**, 3613 (1996). L.P. Kouwenhoven, et.al., Science **278**, 1788 (1997).

[3] *Semiconductor Spintronics and Quantum Computation*, edited by D. D. Awschalom, N. Samarth, and D. Loss (Springer-Verlag, Gerlin, 2002).

[4] D.G. Austing, et.al., Physica B, **249-251**, 206 (1998).

[5] K. Ono, et.al., Science **297**, 1313 (2002).

[6] The second lowest orbital lie $\sim 5\text{meV}$ above the lowest state in our device and the occupation of a (0,2) triplet state can be neglected at small V_S ($< 5\text{mV}$) in low temperatures ($< 2\text{K}$).

[7] S. I. Erlingsson, et.al., Phys. Rev. B **64**, 195306 (2001).

[8] See for example, M. Dobers, et.al., Phys. Rev. Lett. **61**, 1650 (1988).




Figure 3.7(a) (continued)